

Errata

Clinical Guide to Nutrition Care in Kidney Disease Third Edition

Note: The following errata only appears in the first printing of *Clinical Guide to Nutrition Care in Kidney Disease*, 3rd Ed. To determine the print run of your book, consult the middle of the copyright page, which is the page directly before the Table of Contents. The last numeral in the countdown starting with 10 indicates the print run of the book. In the example below, the book would be in its first printing since the 1 is the last digit in the string of numerals.

10 9 8 7 6 5 4 3 2 1

Page and location	Original	Corrected
5, last sentence in first column	The text originally stated that albuminuria is classified as either moderately increased albuminuria with urine albumin levels between 30 mg/d and 300 g/d.	The corrected text should read as follows: between 30 mg/d and 300 mg /d
69, Box 5.2, Vitamin A row	The box originally stated high vitamin A levels can cause anemia and abnormal lipid and metabolism (hypercalcemia).	The corrected text should read as follows: High vitamin A levels can cause anemia and abnormal lipid and calcium metabolism (hypercalcemia).
75, Table 5.1, Vitamin E row	The table originally stated the standard Dietary Reference Intake for vitamin E is 15 IU.	The corrected text should read as follows: 15 mg.
75, Table 5.1, Vitamin E row	The table originally stated the Dietary Reference Intakes for patients on hemodialysis is 400-800 IU.	The updated text reads: 180-360 mg (400-800 IU) with the footnote: 1 IU of the synthetic form (commonly found in supplements) is equivalent to 0.45 mg of alpha-tocopherol.
101, Box 7.9 entry	The box originally stated that the recommended nutrient intake for potassium is 2 to 4 mg/d.	The corrected text should read as follows: 2,000 to 4,000 mg/d
113, Box 8.1, Slowness section	The box originally stated the following: 4-meter walk test. Criterion • 7 seconds or more for people of normal stature scores 1 point • 6 seconds or more for short statured (≤ 173 cm for males and ≤ 159 cm for females) scores 1 point	The corrected text should read as follows: 15-foot walk test. Criterion • 6 seconds or more for people of normal stature scores 1 point • 7 seconds or more for short statured (≤ 173 cm for males and ≤ 159 cm for females) scores 1 point

Errata

Clinical Guide to Nutrition Care in Kidney Disease Third Edition

148, second column, second sentence of full paragraph	The text originally stated that the target blood pressure for adults should be less than 13/80 mm Hg in patients with CKD, with or without DM.	The corrected text should read as follows: 130/80 mm Hg																																								
186, Figure 10.1	The righthand side of the figure originally stated: ↑ phosphate reabsorption	The corrected text should read: ↓ phosphate reabsorption																																								
267, equation at the bottom of the page	The text originally stated <u>that</u> nPCR can be calculated in children as follows: nPCR = 5.43 × Est G V1+ 0.17	The correct text should read as follows: nPCR = 5.43 × Est G + 0.17 V1																																								
328, Box 16.1 entry	The box originally stated that the recommended infusion rate for lipids is less than 0.11 g/kg/min.	The corrected text should read as follows: less than 0.11 g/kg/h																																								
328, Box 16.1 entry, Minerals section	The box originally stated that the sodium recommendations are 1 to 2 mEq/kg/d standard peripheral parenteral nutrition.	The corrected text should read as follows: 1 to 2 mEq/kg/d standard parenteral nutrition .																																								
348, Table 16.6	The table originally stated the following: <table><tr><th></th><th>Amino acids (g)</th><th>Amino acids absorbed (g)</th><th>Dextrose (g)</th></tr><tr><td colspan="4">IPN APD option: 6-L bag compounded using 20% amino acid</td></tr><tr><td>1% Amino acid</td><td>45</td><td>63</td><td>91.5</td></tr><tr><td>1.5% Amino acid</td><td>16</td><td>35</td><td>125</td></tr><tr><td>2% Amino acid</td><td>10</td><td>14 EAA</td><td>100</td></tr></table>		Amino acids (g)	Amino acids absorbed (g)	Dextrose (g)	IPN APD option: 6-L bag compounded using 20% amino acid				1% Amino acid	45	63	91.5	1.5% Amino acid	16	35	125	2% Amino acid	10	14 EAA	100	The corrected text should read as follows: <table><tr><th></th><th>Amino acids (g)</th><th>Amino acids absorbed (g)</th><th>Dextrose (g)</th></tr><tr><td colspan="4">IPN APD option: 6-L bag compounded using 20% amino acid</td></tr><tr><td>1% Amino acid</td><td>60</td><td>30</td><td>85.8</td></tr><tr><td>1.5% Amino acid</td><td>90</td><td>45</td><td>83.4</td></tr><tr><td>2% Amino acid</td><td>120</td><td>60</td><td>81</td></tr></table>		Amino acids (g)	Amino acids absorbed (g)	Dextrose (g)	IPN APD option: 6-L bag compounded using 20% amino acid				1% Amino acid	60	30	85.8	1.5% Amino acid	90	45	83.4	2% Amino acid	120	60	81
	Amino acids (g)	Amino acids absorbed (g)	Dextrose (g)																																							
IPN APD option: 6-L bag compounded using 20% amino acid																																										
1% Amino acid	45	63	91.5																																							
1.5% Amino acid	16	35	125																																							
2% Amino acid	10	14 EAA	100																																							
	Amino acids (g)	Amino acids absorbed (g)	Dextrose (g)																																							
IPN APD option: 6-L bag compounded using 20% amino acid																																										
1% Amino acid	60	30	85.8																																							
1.5% Amino acid	90	45	83.4																																							
2% Amino acid	120	60	81																																							
382, Table 17.1, Vitamin E row	The table originally stated the unit for measuring vitamin E as mcg.	The corrected text should read as follows: mg																																								
383, Box 17.2, Vitamin E row	The box originally stated the unit for measuring vitamin E as mcg.	The corrected text should read as follows: mg																																								