

Errata

Clinical Guide to Nutrition Care in Kidney Disease Third Edition

Note: The following errata only appears in the **first** printing of *Clinical Guide to Nutrition Care in Kidney Disease*, 3rd Ed. To determine the print run of your book, consult the middle of the copyright page, which is the page directly before the Table of Contents. The last numeral in the countdown starting with 10 indicates the print run of the book. In the example below, the book would be in its first printing since the *1* is the last digit in the string of numerals.

10 9 8 7 6 5 4 3 2 1

| Page and location | Original | Corrected | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|--|--|--|-----------------|--------------------------|--------------|--|--|--|--|---------------|----|----|------|-----------------|----|----|-----|---------------|----|--------|-----|---|--|-----------------|--------------------------|--------------|--|--|--|--|---------------|----|----|------|-----------------|----|----|------|---------------|-----|----|----|
| 5, last sentence in first column | The text originally stated that albuminuria is classified as either moderately increased albuminuria with urine albumin levels between 30 mg/d and 300 g/d. | The corrected text should read as follows: between 30 mg/d and 300 mg/d | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 101, Box 7.9 entry | The box originally stated that the recommended nutrient intake for potassium is 2 to 4 mg/d. | The corrected text should read as follows: 2,000 to 4,000 mg/d | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 148, second column, second sentence of full paragraph | The text originally stated that the target blood pressure for adults should be less than 13/80 mm Hg in patients with CKD, with or without DM. | The corrected text should read as follows: 130/80 mm Hg | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 328, Box 16.1 entry | The box originally stated that the recommended infusion rate for lipids is less than 0.11 g/kg/min. | The corrected text should read as follows: less than 0.11 g/kg/h | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 348, Table 16.6 | <p>The table originally stated the following:</p> <table border="1"> <thead> <tr> <th></th> <th>Amino acids (g)</th> <th>Amino acids absorbed (g)</th> <th>Dextrose (g)</th> </tr> </thead> <tbody> <tr> <td colspan="4">IPN APD option: 6-L bag compounded using 20% amino acid</td> </tr> <tr> <td>1% Amino acid</td> <td>45</td> <td>63</td> <td>91.5</td> </tr> <tr> <td>1.5% Amino acid</td> <td>16</td> <td>35</td> <td>125</td> </tr> <tr> <td>2% Amino acid</td> <td>10</td> <td>14 EAA</td> <td>100</td> </tr> </tbody> </table> | | Amino acids (g) | Amino acids absorbed (g) | Dextrose (g) | IPN APD option: 6-L bag compounded using 20% amino acid | | | | 1% Amino acid | 45 | 63 | 91.5 | 1.5% Amino acid | 16 | 35 | 125 | 2% Amino acid | 10 | 14 EAA | 100 | <p>The corrected text should read as follows:</p> <table border="1"> <thead> <tr> <th></th> <th>Amino acids (g)</th> <th>Amino acids absorbed (g)</th> <th>Dextrose (g)</th> </tr> </thead> <tbody> <tr> <td colspan="4">IPN APD option: 6-L bag compounded using 20% amino acid</td> </tr> <tr> <td>1% Amino acid</td> <td>60</td> <td>30</td> <td>85.8</td> </tr> <tr> <td>1.5% Amino acid</td> <td>90</td> <td>45</td> <td>83.4</td> </tr> <tr> <td>2% Amino acid</td> <td>120</td> <td>60</td> <td>81</td> </tr> </tbody> </table> | | Amino acids (g) | Amino acids absorbed (g) | Dextrose (g) | IPN APD option: 6-L bag compounded using 20% amino acid | | | | 1% Amino acid | 60 | 30 | 85.8 | 1.5% Amino acid | 90 | 45 | 83.4 | 2% Amino acid | 120 | 60 | 81 |
| | Amino acids (g) | Amino acids absorbed (g) | Dextrose (g) | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| IPN APD option: 6-L bag compounded using 20% amino acid | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 1% Amino acid | 45 | 63 | 91.5 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 1.5% Amino acid | 16 | 35 | 125 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 2% Amino acid | 10 | 14 EAA | 100 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | Amino acids (g) | Amino acids absorbed (g) | Dextrose (g) | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| IPN APD option: 6-L bag compounded using 20% amino acid | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 1% Amino acid | 60 | 30 | 85.8 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 1.5% Amino acid | 90 | 45 | 83.4 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 2% Amino acid | 120 | 60 | 81 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

Please contact publications@eatright.org with any questions or concerns.