

UNCORRECTED PROOFS

Food & Supplement Medication Interactions

Nutrition and Dietetics
Educators and Preceptors

EDITORS:

Lauren Roberson, PhD, RDN, LD

Avani Patel, PharmD, RDN, MDN, LD

**eat
right.**® Academy of
Nutrition
and Dietetics

UNCORRECTED PROOFS

Contents

Contributors	ii
Reviewers	iii
Common Abbreviations	iv
Preface	vii
Introduction	1
How to Use This Guide	1
Defining Food and Supplement Medication Interactions	2
Assessing Medications as Part of the Nutrition Care Process.	4
Understanding Drug Metabolism	5
Drugs by Generic Name	9
Bibliography.	xx

SAMPLE
Not for Print
or Sale

UNCORRECTED PROOFS

Introduction

This resource provides notable and documented interactions between food, including some dietary supplements, and medications (also referred to as drugs) that nutrition practitioners are likely to encounter in a clinical setting. The information provided is not all-inclusive or exhaustive as the availability of medications is constantly changing and some have limited or no effects on nutrition status. Additionally, research on adverse effects of medications impacting nutrition status is evolving. Information presented here is reflective of the research available at the time of publication. Other sources may need to be consulted for some medications and for dietary supplements. This resource is intended to be used in conjunction with other clinical practice resources.

How to Use This Guide

Over 560 medications are included and listed alphabetically by generic name. The following information is presented as it applies to each medication:

- **DRUG NAME:** Generic name of drug currently available in the US
- **DRUG CLASS:** Category (or categories) in which drugs are classified according to their mechanism of action and intended effect
- **COMMON USES:** Conditions for which the drug is most commonly prescribed
- **DIET CONSIDERATIONS:** Information related to timing of taking medications in relation to meals and whether intake of specific substances (eg, grapefruit products, caffeine, alcohol), and/or specific macro- or micronutrients may need to be adjusted, limited, or avoided.
- **FOOD (OR NUTRIENT) DRUG INTERACTIONS:** Brief explanation of how specific foods, ingredients, or macro- or micronutrients can either increase or decrease drug effects (and side effects), or how drugs may impact macro- or micronutrient status (eg, some drugs can increase micronutrient loss).
- **DRUG-SUPPLEMENT INTERACTIONS:** Information, as available, related to dietary and herbal supplements ranging from food-based (eg, garlic and maitake) to vitamin and mineral supplements to substances used recreationally or for medical purposes (eg, cannabis, products containing cannabinoids, mushrooms, and melatonin), and their effects on increasing or decreasing a drug's effects. Note that cannabis listed in this resource refers to the oral form and it is considered a controlled substance and not a dietary supplement.

Information related to supplements is included only if there was sufficient evidence to document the interaction, such as in UpToDate. Due to limitations in published research on dietary supplements and variability in the composition of dietary supplements, clinical data is often insufficient to make conclusive judgments about the incidence and severity of potential drug-supplement interactions. The registered dietitian nutritionist (RDN) should consult available resources on specific dietary supplements and discuss findings with the health

UNCORRECTED PROOFS

care team to make informed decisions regarding use of dietary supplements in conjunction with certain medications.

- **NUTRITION-RELATED SIDE EFFECTS:** Includes the most common side effects that influence nutrition status observed in adult patients (unless otherwise noted) taking each drug. This includes side effects related to gastrointestinal function, appetite, weight, hydration status, laboratory tests, and blood levels of specific vitamins or minerals. Percentages associated with various side effects are provided for some drugs; these reflect documented prevalence of the side effect (based on available data from UpToDate and peer-reviewed studies).
- **MONITOR:** Lab tests that should be monitored as well as anthropometric data, vitals (eg, blood pressure), and related signs and symptoms that should be monitored by the RDN in the treatment of patients who have been prescribed the given drug.

All information has been gathered from peer-reviewed sources, which may include professional databases, articles, book chapters, or website content. Refer to the bibliography at the back of this resource for more details.

NOTE: The term drug can refer to any chemical substance that affects the functioning of the body, typically used for therapeutic, diagnostic, or recreational purposes, whereas medication specifically refers to a drug that is used for therapeutic purposes to prevent, treat, or manage medical conditions or symptoms. Thus, all medications are considered drugs, while not all drugs are medications. Throughout this resource, both drug and medication terminology are used interchangeably to describe any substance being used for therapeutic or diagnostic purposes and under the supervision of a health care professional.

Defining Food and Supplement Medication Interactions

Foods (including ingredients and nutrients) and supplements can enhance, delay, or decrease the absorption of some drugs, and drugs can have a wide range of effects on food intake, nutrient metabolism, and nutritional status. Consequently, an interaction occurs when a food (or nutrient) affects the metabolism or absorption of a drug, or when a drug affects the absorption, metabolism, or excretion of a nutrient. These interactions can alter the effectiveness of a medication, leading to increased risk for toxicity or significantly reduced therapeutic effect.^{1,2} Drug-food/nutrient interactions can also affect nutrient intake, absorption, and metabolism and the overall nutritional health of an individual.

UNCORRECTED PROOFS

Abacavir

Antiretroviral, nucleotide reverse transcriptase inhibitor (NRTI)

COMMON USES HIV infection

DIET CONSIDERATIONS Take with or without food.

Oral solution contains sorbitol; consider other dietary sources of sugar alcohols to prevent GI side effects.

NUTRITION-RELATED SIDE EFFECTS

- Nausea, 7%–19%
- Diarrhea, 7%
- Abdominal pain, <6%
- Gastritis, <6%
- Elevated serum amylase, 2%–4%
- Vomiting, 2%
- Hepatomegaly with steatosis
- Lactic acidosis
- Mild hyperglycemia (more common in pediatric pts)

Abdominal discomfort or diarrhea may occur in pts who are sensitive to fructose.

MONITOR *Lab tests:* CBC, fasting lipid panel, liver function (AST, ALT)

Abatacept

Selective T-cell costimulation blocker, antirheumatic (disease modifying)

COMMON USES aGVHD prophylaxis, psoriatic arthritis, rheumatoid arthritis

DRUG-FOOD (OR NUTRIENT) INTERACTIONS Powder for injection may contain maltose; may result in falsely ↑ BG on day of infusion.

NUTRITION-RELATED SIDE EFFECTS

- Hypermagnesemia, 18% when prescribed for aGVHD
- Nausea, ≥10% when prescribed for rheumatoid arthritis; severe nausea in 5% when prescribed for aGVHD
- Diarrhea, severe, 6% when prescribed for aGVHD
- Dyspepsia, 6%
- Abdominal pain
- Severe diverticulitis

MONITOR *Lab tests:* Mg, kidney function (Cr, GFR), anemia (Hgb, Hct)

Other: GI symptoms, BP, temperature in pts with aGVHD

Acamprosate

GABA agonist/glutamate antagonist

COMMON USES Moderate to severe alcohol use disorder

DIET CONSIDERATIONS Take with or without food. Avoid alcohol.

DRUG-FOOD (OR NUTRIENT) INTERACTIONS Alcohol ↓ drug efficacy.

NUTRITION-RELATED SIDE EFFECTS

- Diarrhea, 17%
- Anorexia, 5%
- Flatulence, 4%
- Nausea, 4%
- Abdominal pain, ≥1%
- ↑ Appetite, ≥1%
- Constipation, ≥1%
- Dysgeusia, ≥1%
- Dyspepsia, ≥1%
- HTN, ≥1%
- Peripheral edema, ≥1%
- Vomiting, ≥1%
- Wt gain, ≥1%
- AKI
- Angioedema

MONITOR *Lab tests:* Monitor for nutrient deficiencies (eg, folate, thiamin, and vitamins A, B6, B12, D, E, and K).

Other: Wt (baseline), kidney function (baseline), fluid and electrolyte status

UNCORRECTED PROOFS

Lomitapide

Antilipemic agent, microsomal triglyceride transfer protein (MTTP) inhibitor

COMMON USES Homozygous familial hypercholesterolemia

DIET CONSIDERATIONS Take capsule with 8 oz of plain water ≥ 2 h after a low-fat meal; low-fat diet recommended ($<20\%$ calories from fat).

Avoid grapefruit products.

Limit or avoid alcohol (≤ 1 alcoholic beverage per day).

DRUG-FOOD (OR NUTRIENT) INTERACTIONS Drug can \downarrow absorption of fat-soluble vitamins and fatty acids; consider supplementation of vitamin E, linoleic acid, α -linolenic acid, eicosapentaenoic acid, and docosahexaenoic acid.

Grapefruit products can \uparrow drug serum concentration.

Alcohol can \uparrow hepatotoxic effect of drug.

DRUG-SUPPLEMENT INTERACTIONS Ginkgo biloba may \uparrow drug serum concentration. Drug may \uparrow serum concentration of red yeast rice.

NUTRITION-RELATED SIDE EFFECTS

- Abdominal discomfort
- Abdominal distention
- Abdominal pain
- Constipation
- Diarrhea
- Dyspepsia
- Flatulence
- Gastroenteritis
- GERD
- N/V
- Wt loss

MONITOR *Lab tests:* liver function (ALT, AST, ALP, bilirubin)

Loperamide

Antidiarrheal

COMMON USES Diarrhea

DIET CONSIDERATIONS *Chewable tablet:* Take only on empty stomach 1 h before or 2 h after a meal. Drink plenty of water, especially if diarrhea persists.

Some formulations may contain Na.

DRUG-FOOD (OR NUTRIENT) INTERACTIONS Caution with children or elderly due to \uparrow risk of dehydration or electrolyte imbalance.

DRUG-SUPPLEMENT INTERACTIONS Kratom may \uparrow CNS depressant effects of drug.

NUTRITION-RELATED SIDE EFFECTS

- Constipation, 2%-5%
- Abdominal cramps, $\leq 3\%$
- Nausea, $\leq 3\%$

MONITOR *Lab tests:* electrolytes, liver function (ALP, ALT, AST)

Other: fluid status

Lopinavir/Ritonavir

Antiretroviral, protease inhibitor (anti-HIV)

COMMON USES HIV-1 infection